

Bibliography: Percutaneous Electrical Nerve Field Stimulation (PENFS)

Clinical Outcomes Studies in Children and Adolescents 8-21 years with Pain Associated DGBIs

1. Waheed U, VonAxelson A, El-Chammas K, Graham K, Mansi S, Hardy J, Miller MM, Sahay R, Santucci NR. Percutaneous Electrical Nerve Field Stimulation Improves Weight and Other Anthropometric Outcomes in Children and Adolescents With Functional Dyspepsia Who Restrict Their Oral Intake. *Neurogastroenterol Motil.* 2026;38(2):e70273.
2. Gordon M, Benninga MA, Borlack R, Borrelli O, Chogle A, Darbari A, Dolinsek J, Groen J, Khlevner J, Di Lorenzo C, Person H, Saps M, Sinopoulou V, Snyder J, Tabbers M, Thapar N, Vlioger A. ESPGHAN/NASPGHAN guidelines for treatment of irritable bowel syndrome and functional abdominal pain-not otherwise specified in children aged 4–18 years. *J Pediatr Gastroenterol Nutr.* 2025; 81: 442-471.
3. Dorfman L, El-Chammas K, Graham K, Sahay R, Hardy J, Kaul A, Santucci N. Repeat round of auricular percutaneous electrical nerve field stimulation for pediatric disorders of gut brain interaction. *J Pediatr Gastroenterol Nutr.* 2025;81:234-245.
4. Santucci NR, Waheed U, Li J, Mansi S, Graham K, Hardy J, Miller MM, Sahay R, El-Chammas K. Auricular Allodynia is Associated With Worse Outcomes in Children With Functional Abdominal Pain Disorders Using Neurostimulation. *Neuromodulation.* 2025;28:840-846.
5. Kolacz J, Roath OK, Lewis GF, Karrento K. Cardiac Vagal Efficiency Is Enhanced by Percutaneous Auricular Neurostimulation in Adolescents With Nausea: Moderation by Antidepressant Drug Exposure. *Neurogastroenterol Motil.* 2025 May;37(5):e15007.
6. Santucci NR, Beigarten AJ, Khalid F, El-Chammas KI, Graham K, Sahay R, Fei L, Rich K, Mellon M. Percutaneous Electrical Nerve Field Stimulation in Children and Adolescents With Functional Dyspepsia- Integrating a Behavioral Intervention. *Neuromodulation.* 2024;27:372-381.
7. Chogle A, El-Chammas K, Santucci N, Grimm M, Dorfman L, Graham K, Kelly DR, Dranove JE, Rosen R, Nurko S, Croffie J, Balakrishnan K, Chiou EH, Zhang L, Simpson P, Karrento K. A multicenter registry study on percutaneous electrical nerve field stimulation for pediatric disorders of gut-brain interaction. *J Pediatr Gastroenterol Nutr.* 2024;78:817-826.
8. Santucci NR, Sahay R, El-Chammas KI, Graham K, Wheatley M, Vandenbrink M, Hardy J, Fei L. Percutaneous electrical nerve field stimulation compared to standard medical therapy in adolescents with functional abdominal pain disorders. *Front Pain Res (Lausanne).* 2023;4:1251932.
9. Chogle A, Visnagra K, Janchoi J, Tran T, Davis R, Callas N and Ornelas E. Prospective study of the effect of auricular percutaneous electrical nerve field stimulation on quality of life in children with pain related disorders of gut-brain interaction. *Front Pain Res (Lausanne).* 2023;4:1223932.
10. Castillo DF, Denson LA, Haslam DB, Hommel KA, Ollberding NJ, Sahay R, Santucci NR. The microbiome in adolescents with irritable bowel syndrome and changes with percutaneous electrical nerve field stimulation. *Neurogastroenterol Motil.* 2023;35(7):e14573.
11. Bora G, Atkinson SN, Pan A, Sood M, Salzman N, Kovacic K. Impact of auricular percutaneous electrical nerve field stimulation on gut microbiome in adolescents with irritable bowel syndrome: A pilot study. *J Dig Dis.* 2023;24(5):348-358.
12. Amornluck Krasaelap, Manu R Sood, B U K Li et al. Efficacy of Auricular Neurostimulation in Adolescents with Irritable Bowel Syndrome in a Randomized, Double-Blind Trial. *Clin Gastroenterol Hepatol.* 2020;18(9):1987-1994.e2.

13. Kovacic K, Kolacz J, Lewis GF, Porges SW. Impaired Vagal Efficiency Predicts Auricular Neurostimulation Response in Adolescent Functional Abdominal Pain Disorders. *Am J Gastroenterol*. 2020;115(9):1534-1538.
14. Kovacic K, Hainsworth K, Sood M, Chelimsky G, Unteutsch R, Nugent M, Simpson P, Miranda A. Neurostimulation for abdominal pain-related functional gastrointestinal disorders in adolescents: a randomised, double-blind, sham-controlled trial. *Lancet Gastroenterol Hepatol*. 2017;2(10):727-737.
15. Roberts A, Sithole A, Sedghi M, Walker CA, Quinn TM. Minimal adverse effects profile following implantation of periauricular percutaneous electrical nerve field stimulators: a retrospective cohort study. *Med Devices (Auckl)*. 2016;9:389-393.

Clinical Outcomes Studies in Children and Adolescents 8-21 years with other DGBIs

16. Karrento K, Venkatesan T, Zhang L, Pawela L, Simpson P, Li BUK. Percutaneous Electrical Nerve Field Stimulation for Drug-Refractory Pediatric Cyclic Vomiting Syndrome. *J Pediatr Gastroenterol Nutr*. 2023;77(3):347-353.
17. Karrento K, Zhang L, Conley W, Qazi Z, Venkatesan T, Simpson P, Li BUK. Percutaneous electrical nerve field stimulation improves comorbidities in children with cyclic vomiting syndrome. *Front Pain Res (Lausanne)*. 2023;4:1203541.

Adult studies, non-DGBI.

18. Woodbury A, Krishnamurthy LC, Bohsali A, Krishnamurthy V, Smith JL, Gebre M, Tyler K, Vernon M, Crosson B, Kalangara JP, Napadow V, Allen JW, Harper D. Percutaneous electric nerve field stimulation alters cortical thickness in a pilot study of veterans with fibromyalgia. *Neurobiol Pain*. 2022;12:100093.
19. Woodbury A, Krishnamurthy V, Gebre M, Napadow V, Bicknese C, Liu M, Lukemire J, Kalangara J, Cui X, Guo Y, Sniecinski R, Crosson B. Feasibility of Auricular Field Stimulation in Fibromyalgia: Evaluation by Functional Magnetic Resonance Imaging, Randomized Trial. *Pain Med*. 2021;22:715-726.

Cost analysis in adolescents with Irritable Bowel Syndrome

20. Shah E, Eswaran S, Harer K, et al. Percutaneous electrical nerve field stimulation for adolescents with irritable bowel syndrome: cost-benefit and cost-minimization analysis. *J Ped Gastro Nutr*. 2024;1-6.

Basic Science Animal Model and Mechanism

21. Babygirija R, Sood M, Kannampalli P, Sengupta JN, Miranda A. Percutaneous electrical nerve field stimulation modulates central pain pathways and attenuates post-inflammatory visceral and somatic hyperalgesia in rats. *Neuroscience*. 2017;356:11-21.